



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
&= \frac{1}{16}(\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2}) + \frac{1}{16}(\sqrt{10 + 4\sqrt{5}} + \sqrt{10 - 4\sqrt{5}} \\
&\quad - \sqrt{30 + 12\sqrt{5}} - \sqrt{30 - 12\sqrt{5}}) \\
&= \frac{1}{16}(\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2}) + \frac{1}{16}(1 - \sqrt{3})(\sqrt{10 + 4\sqrt{5}} + \sqrt{10 - 4\sqrt{5}}).
\end{aligned}$$

By simplifying the expression in the last parenthesis and then collecting under the radical sign, we have, after a simple reduction,

$$\sin 3^\circ = \frac{1}{16}(\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2}) + \frac{1}{16}(\sqrt{5 + \sqrt{5}} - \sqrt{15 + 3\sqrt{5}}).$$

Also solved by L. E. MENSENKAMP, J. L. RILEY, HORACE OLSON, H. S. UHLER, H. C. FEEMSTER, FLORENCE RAE, R. M. MATHEWS, H. L. AGARD, GILBERT A. AULT, FRANK IRWIN, C. E. GITHENS, A. M. HARDING, and the PROPOSER.

491 (Algebra). Proposed by J. W. LASLEY, University of North Carolina.

Solve the equations $xy = x^2 - y^2$ and $x^2 + y^2 = x^3 - y^3$ for x and y .

SOLUTION BY J. W. BALDWIN, Ann Arbor, Michigan.

Solving $xy = x^2 - y^2$ for x in terms of y we have $x = \frac{1}{2}(1 + \sqrt{5})y$ and $x = \frac{1}{2}(1 - \sqrt{5})y$. These values substituted in $x^2 + y^2 = x^3 - y^3$ give, after simplification, $y^2(y - \frac{1}{2}\sqrt{5}) = 0$ (1) and $y^2(y + \frac{1}{2}\sqrt{5}) = 0$ (2). From (1), $y = 0, 0, \frac{1}{2}\sqrt{5}$ and from (2) $y = 0, 0, -\frac{1}{2}\sqrt{5}$. Hence, the corresponding values of x are $0, 0, (5 + \sqrt{5})/4$ and $0, 0, (5 - \sqrt{5})/4$. From (1) or (2) it is seen that two branches of the curve represented by the second equation pass through the origin. It is readily determined that these branches are imaginary and, hence, the origin is a conjugate point.

Also solved by S. E. RASOR, F. H. HOLESTIN, ELGIN E. GROSECLOSE, H. N. CARLETON, A. M. HARDING, J. L. RILEY, POLYCARP HANSEN, E. B. ESCOTT, G. Y. SOSNOW, J. Q. McNATT, O. S. ADAMS, HORACE OLSON, T. C. AMICK, L. E. LUNN, PAULINE SPERRY and the PROPOSER.

433 (Calculus). Proposed by LOUIS O'SHAUGHNESSY, University of Pennsylvania.

Solve the differential equation

$$\frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} - \frac{y}{x} = 0.$$

I. SOLUTION BY EMIL L. POST, New York City.

We have

$$x \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} = y. \quad (1)$$

Operating on both sides by $d^{\frac{1}{2}}/dx^{\frac{1}{2}}$ (see next problem), we have

$$\frac{d^{\frac{1}{2}}}{dx^{\frac{1}{2}}} \left[x \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} \right] = \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}}. \quad (2)$$

But for any operation $f(D)$

$$f(D)u \cdot v = uf(D)v + \frac{du}{dx} \frac{f'(D)v}{1!} + \frac{d^2u}{dx^2} \frac{f''(D)v}{2!} + \dots$$

Let

$$f(D) = D^{\frac{1}{2}} = \frac{d^{\frac{1}{2}}}{dx^{\frac{1}{2}}}; \quad u = x; \quad v = \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}}.$$

Then

$$\frac{d^{\frac{1}{2}}}{dx^{\frac{1}{2}}} \left[x \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} \right] = x \frac{d^{\frac{1}{2}}}{dx^{\frac{1}{2}}} \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} + \frac{1}{2} \frac{d^{-\frac{1}{2}}}{dx^{-\frac{1}{2}}} \frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} = x \frac{dy}{dx} + \frac{1}{2}y. \quad (3)$$

Substituting from the original equation and (3) into (2), we have

$$x \frac{dy}{dx} + \frac{1}{2}y = \frac{y}{x}. \quad (4)$$

From this we have

$$\log y = -1/x - \frac{1}{2} \log x + \log C,$$

or

$$y = Ce^{-(1/x)}x^{-\frac{1}{2}}.$$

II. SOLUTION BY THE PROPOSER.

According to Professor Kelland (*Trans. Royal Society of Edinburgh*, Vols. XIV and XVI) the general differential operator may be defined as follows:

$$\frac{d^\mu x^n}{dx^\mu} = (-1)^\mu \frac{\Gamma(-n + \mu)}{\Gamma(-n)} x^{n-\mu},$$

for all values of n and μ . We have $\Gamma(n+1) = n\Gamma(n)$.

Let us assume $y = A_0 + A_1x^{-\frac{1}{2}} + A_2x^{-1} + A_3x^{-\frac{3}{2}} + \dots$.

Then

$$\frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} = (-1)^{\frac{1}{2}} \left\{ \frac{\Gamma(1)}{\Gamma(\frac{1}{2})} A_1x^{-1} + \frac{\Gamma(\frac{3}{2})}{\Gamma(1)} A_2x^{-\frac{3}{2}} + \frac{\Gamma(2)}{\Gamma(\frac{3}{2})} A_3x^{-2} + \dots \right\}$$

and

$$\frac{y}{x} = A_0x^{-1} + A_1x^{-\frac{3}{2}} + A_2x^{-2} + A_3x^{-\frac{5}{2}} + \dots$$

Hence,

$$i \frac{\Gamma(1)}{\Gamma(\frac{1}{2})} A_1 = A_0 \text{ and } A_1 = \frac{\sqrt{\pi}}{i} A_0 = -i\sqrt{\pi}A_0.$$

For, since $\Gamma(p)\Gamma(1-p) = \pi/\sin p\pi$ when p is a fraction less than one, then $\Gamma(\frac{1}{2}) = \sqrt{\pi}$. Also, $\Gamma(1) = 1$.

Furthermore,

$$i \frac{\Gamma(\frac{3}{2})}{\Gamma(1)} A_2 = A_1 = -i\sqrt{\pi}A_0, \text{ or } A_2 = -2A_0,$$

since $\Gamma(\frac{3}{2}) = \frac{1}{2}\Gamma(\frac{1}{2}) = \frac{1}{2}\sqrt{\pi}$; also

$$i \frac{\Gamma(2)}{\Gamma(\frac{3}{2})} A_3 = A_2 = -2A_0, \text{ or } A_3 = -\frac{\sqrt{\pi}}{i} A_0 = i\sqrt{\pi}A_0.$$

Finally, $y = A_0(1 - i\sqrt{\pi}x^{-\frac{1}{2}} - 2x^{-1} + i\sqrt{\pi}x^{-\frac{3}{2}} + \dots)$.

Note.—This problem, which is very similar to one solved by Professor Kelland, was submitted because it was thought that it might prove of interest to the readers of the MONTHLY. The proposer would be glad to see some discussion of the subject of general differentiation.

The above solution may be considered as a reply to the following question:

360 (Calculus.) Proposed by **ELMER SCHUYLER**, Brooklyn, New York.

What interpretation must be given to

$$\frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} \text{ so that } \frac{d^{\frac{1}{2}}}{dx^{\frac{1}{2}}} \left(\frac{d^{\frac{1}{2}}y}{dx^{\frac{1}{2}}} \right) = \frac{dy}{dx}?$$

435 (Calculus.) Proposed by **B. F. FINKEL**, Drury College.

Show that

$$\int_0^\infty e^{-x^2 - a^2/x^2} dx = \frac{\sqrt{\pi}}{2e^{2a}}$$

by a transformation, rather than by the usual method of differentiating under the sign of integration, as, for example, in Byerly's *Integral Calculus*, page 106-107.